Petrushyn Dmytro, Cand. of Sciences Physical Education and Sports, Dnipro State University of Internal Affairs, Dnipro, Ukraine, https://orcid.org/0000-0001-9207-0804

CARDIAC STRUCTURAL ADAPTATIONS IN ELITE BOXERS: CLINICAL IMPLICATIONS AND DIFFERENTIATION FROM PATHOLOGY

Abstract

Background: Boxing represents a unique combination of aerobic and anaerobic exercise that induces specific cardiac adaptations. Understanding these physiological changes is crucial for differentiating the athlete's heart from pathological conditions. **Objective:** To analyze cardiac structural adaptations in elite boxers and establish clinical criteria for differentiating physiological remodeling from pathological conditions such as hypertrophic cardiomyopathy. **Methods:** A comprehensive literature review was conducted examining cardiac morphology, electrocardiographic patterns, and cardiovascular magnetic resonance findings in elite boxers. Studies analyzing both male and female boxers across various competitive levels were included. **Results:** Elite boxers demonstrate characteristic cardiac adaptations including increased left ventricular wall thickness, chamber enlargement, and specific electrocardiographic patterns. These adaptations correlate with training intensity and duration but remain within physiological limits. Gender-specific differences in cardiac remodeling patterns were identified, with male boxers showing more pronounced structural changes. Autonomic modulation, assessed through heart rate variability, provides additional markers for distinguishing physiological from pathological adaptations. **Conclusions:** Cardiac adaptations in elite boxers represent physiological responses to training demands. Comprehensive evaluation using established

criteria, including cardiovascular magnetic resonance and heart rate variability analysis, enables accurate differentiation from pathological conditions. Regular monitoring is essential for athlete safety and performance optimization.

Keywords: boxing, athlete's heart, cardiac remodeling, hypertrophic cardiomyopathy, heart rate variability, cardiovascular magnetic resonance, sports cardiology.

Introduction

Boxing is a high-intensity combat sport that demands exceptional physical capabilities, combining explosive power, speed, endurance, and technical proficiency. The physiological demands of boxing training and competition induce specific adaptations in multiple body systems, with cardiovascular adaptations being among the most pronounced and clinically significant (Chaabene et al., 2015).

The «athlete's heart» phenomenon, characterized by structural and functional cardiac adaptations to chronic exercise training, has been extensively documented across various sports disciplines. However, boxing presents unique challenges due to its distinct training profile that combines high-intensity interval work, sustained aerobic conditioning, and explosive anaerobic efforts (Davis et al., 2015). Understanding these adaptations is crucial for sports medicine practitioners, as distinguishing physiological remodeling from pathological conditions such as hypertrophic cardiomyopathy (HCM) remains a critical clinical challenge.

Elite boxers undergo rigorous training regimens that typically include cardiovascular conditioning, technical skill development, strength training, and sport-specific sparring sessions. These training modalities create a mixed physiological stimulus that differs substantially from purely endurance or resistance-based sports (Chaabene et al., 2015). The cardiac response to such training includes chamber enlargement, increased wall thickness, and alterations in

autonomic regulation that must be carefully evaluated to ensure athlete safety (Ghani & Sharma, 2020).

Recent advances in cardiac imaging, particularly cardiovascular magnetic resonance (CMR), and the refinement of electrocardiographic interpretation criteria specifically designed for athletes have improved our ability to differentiate physiological from pathological cardiac adaptations (Drezner et al., 2017). Additionally, heart rate variability (HRV) analysis has emerged as a valuable non-invasive tool for assessing autonomic function and training adaptation in athletes (Kiss et al., 2016).

This article aims to comprehensively analyze the cardiac structural adaptations observed in elite boxers, examine the clinical implications of these changes, and provide evidence-based guidance for distinguishing physiological remodeling from pathological conditions that may predispose athletes to adverse cardiac events.

Boxing competition and training impose substantial cardiovascular demands characterized by repeated high-intensity efforts interspersed with brief recovery periods. During competitive bouts, heart rates can reach 85-95% of maximum, with sustained elevations throughout the contest (Davis et al., 2015). The activity profile of elite male amateur boxers demonstrates that approximately 33% of bout time involves high-intensity activity, creating significant cardiac workload.

The training regimen of elite boxers typically encompasses multiple daily sessions including roadwork for aerobic conditioning, technical sparring, pad work for explosive power development, and strength conditioning. This multifaceted approach creates a combined volume and pressure overload on the heart, stimulating both eccentric and concentric hypertrophy patterns (Chaabene et al., 2015).

Research examining cardio-respiratory responses during simulated boxing contests in elite-level boxers reveals significant cardiovascular stress, with

substantial oxygen consumption and ventilatory demands throughout three-minute rounds (Kamandulis et al., 2018). These physiological stressors, applied consistently over years of training, drive the adaptive remodeling observed in the athlete's heart.

Understanding these demands is essential for interpreting the structural and functional cardiac changes observed in boxers, as the magnitude and pattern of adaptation directly correlate with the specific training stimulus applied.

The most prominent cardiac adaptation in elite boxers involves left ventricular (LV) remodeling. Years of intensive training induce increases in LV wall thickness, chamber dimensions, and mass that reflect the heart's response to sustained workload. Studies utilizing CMR imaging have established reference ranges for cardiac structure in athletes, demonstrating that boxers typically fall within the upper limits of physiological adaptation (Petersen et al., 2017).

The pattern of LV hypertrophy in boxers tends to be balanced, reflecting the mixed nature of their training stimulus. Unlike pure endurance athletes who develop predominantly eccentric hypertrophy (chamber enlargement with proportional wall thickening) or pure strength athletes who may develop more concentric patterns (wall thickening with less chamber enlargement), boxers demonstrate characteristics of both patterns (Utomi et al., 2013).

Meta-analyses examining the morphology and function of the male athlete's heart across different training modalities confirm that combat sports athletes, including boxers, develop significant but physiological increases in LV mass, with typical wall thickness measurements ranging from 11-13 mm (Utomi et al., 2013). These values generally remain below the pathological threshold of 15 mm used to diagnose HCM, though exceptional cases may approach or slightly exceed this value while remaining physiological.

Beyond wall thickness, chamber dimensions provide crucial information about cardiac adaptation. Elite boxers typically demonstrate enlarged LV end-

diastolic volumes compared to sedentary individuals, reflecting increased stroke volume capacity. This enlargement occurs proportionally with wall thickening, maintaining normal relative wall thickness ratios that distinguish physiological from pathological remodeling (Petersen et al., 2017).

Right ventricular adaptation, though less extensively studied than left ventricular changes, also occurs in response to boxing training. Standardized CMR protocols have enabled better characterization of biventricular remodeling patterns in athletes (Kramer et al., 2008).

Left atrial enlargement represents another common finding in endurance-trained athletes and is observed in elite boxers. This adaptation facilitates enhanced diastolic filling and contributes to increased stroke volume. However, significant atrial enlargement requires careful evaluation to exclude underlying pathology or increased risk of atrial arrhythmias.

Emerging evidence highlights important gender differences in cardiac adaptation to athletic training. Studies examining male and female Olympic boxers reveal that while both sexes develop cardiac adaptations, males typically demonstrate more pronounced structural changes (Bianco et al., 2023).

Research investigating gender dimorphism in combat sports athletes, including boxers and acrobats, indicates that physiological adaptations, including cardiac remodeling, follow sex-specific patterns related to hormonal influences, body composition, and training responses (Bachynska et al., 2023, 2025). Female athletes generally develop less pronounced LV hypertrophy and chamber enlargement compared to male counterparts training at similar intensities, though they may demonstrate relatively greater increases in contractile function.

These gender differences have important clinical implications for establishing diagnostic criteria and reference ranges. Application of universal cutoff values without consideration of sex-specific patterns may lead to either overdiagnosis of pathology in male athletes or underdiagnosis in female athletes.

The electrocardiogram (ECG) provides essential information about cardiac electrical activity and is a cornerstone of pre-participation cardiovascular screening in athletes. Elite boxers commonly exhibit ECG changes that reflect underlying structural adaptations and enhanced vagal tone.

Training-related ECG changes in boxers include sinus bradycardia, first-degree atrioventricular block, incomplete right bundle branch block, early repolarization patterns, and increased QRS voltage suggestive of LV hypertrophy (Drezner et al., 2017). These findings are generally considered benign adaptations to training and do not warrant additional investigation in asymptomatic athletes with normal physical examination findings.

International consensus criteria for ECG interpretation in athletes provide standardized approaches to distinguishing normal training-related findings from abnormal patterns that require further evaluation (Drezner et al., 2017). Studies comparing different ECG interpretation criteria demonstrate that athlete-specific criteria reduce false-positive rates while maintaining sensitivity for detecting underlying cardiac pathology (Riding et al., 2015).

Certain ECG findings raise concern for underlying cardiac disease and warrant comprehensive evaluation regardless of symptoms. These include T-wave inversions beyond V1, pathological Q waves, left bundle branch block, QTc prolongation or shortening, profound sinus bradycardia below 30 beats per minute, and ventricular pre-excitation patterns (Drezner et al., 2017).

In boxers presenting with these abnormal patterns, further investigation with echocardiography, CMR, and possibly genetic testing is essential to exclude conditions such as HCM, arrhythmogenic right ventricular cardiomyopathy, long QT syndrome, or other inherited cardiac disorders.

Heart rate variability represents beat-to-beat variations in heart rate and reflects autonomic nervous system regulation of cardiac function. HRV analysis

has emerged as a valuable tool for assessing training adaptation, recovery status, and autonomic balance in athletes (Morgan & Mora, 2017).

Studies examining HRV in combat sports athletes, including boxers, demonstrate characteristic patterns associated with high-level training. Elite athletes typically exhibit elevated HRV compared to untrained individuals, reflecting enhanced parasympathetic tone and cardiac autonomic regulation (Kiss et al., 2016). This adaptation contributes to resting bradycardia and represents a favorable training response.

Research investigating HRV in acrobats and combat athletes across long-term training periods, with consideration of gender differences, reveals that autonomic adaptations parallel structural cardiac changes (Bachynska et al., 2023). Regular HRV monitoring provides insights into training load tolerance, recovery adequacy, and potential overtraining states.

Day-to-day HRV monitoring in elite athletes enables tracking of training responses and early detection of maladaptation (Plews et al., 2016). In professional athletes across various sports, including soccer players and swimmers, HRV metrics correlate with performance capacity and injury risk (Lima-Borges et al., 2018; Rabbani et al., 2019).

For boxers specifically, HRV biofeedback and monitoring may optimize training prescription and recovery strategies, potentially enhancing performance while reducing injury and illness risk (Morgan & Mora, 2017). During intensive training periods or overload microcycles, HRV monitoring helps distinguish productive training stress from excessive fatigue (Schneider et al., 2019).

Additionally, HRV assessment following specific training stimuli, such as high-intensity interval sessions or environmental stressors like sauna exposure, provides insights into individual recovery patterns and adaptation capacity (Leicht et al., 2018).

Distinguishing physiological cardiac hypertrophy in elite boxers from pathological conditions, particularly HCM, represents a critical clinical challenge with profound implications for athlete safety and competitive participation. HCM remains the leading cause of sudden cardiac death in young athletes, making accurate diagnosis essential (Harmon et al., 2015).

Several morphological features help differentiate the athlete's heart from HCM. Physiological hypertrophy in boxers typically demonstrates symmetric or mildly asymmetric LV wall thickening with proportional chamber enlargement, maintaining normal LV cavity size (typically >55 mm end-diastolic dimension). In contrast, HCM often presents with marked asymmetric septal hypertrophy, small or normal LV cavity dimensions, and systolic anterior motion of the mitral valve (Elliott et al., 2014).

Wall thickness measurements exceeding 15 mm raise concern for HCM, though occasional elite athletes may demonstrate physiological wall thickness approaching or slightly exceeding this threshold. In such cases, additional evaluation is mandatory to exclude pathology (Elliott et al., 2014).

Diastolic function assessment provides valuable discriminatory information. Athletes with physiological hypertrophy maintain normal or supranormal diastolic function, while HCM patients typically demonstrate impaired relaxation and elevated filling pressures. CMR evaluation of diastolic parameters, tissue characterization, and the presence of late gadolinium enhancement (suggesting myocardial fibrosis) helps distinguish physiological from pathological remodeling (Kramer et al., 2008).

Systolic function in athletic hypertrophy remains normal or enhanced, whereas HCM may demonstrate hyperdynamic systolic function with LV outflow tract obstruction or, in advanced cases, systolic dysfunction.

A classic approach to resolving diagnostic uncertainty involves a period of detraining. Physiological cardiac hypertrophy partially regresses with cessation of

training over 3-6 months, whereas pathological hypertrophy in HCM remains unchanged or progresses. While valuable, this approach delays competitive participation and creates psychological stress for athletes, making it less practical for elite competitors at the peak of their careers.

Comprehensive evaluation includes detailed family history assessment for premature sudden cardiac death, HCM, or other inherited cardiac conditions. In cases of diagnostic uncertainty, genetic testing for HCM-associated sarcomeric protein mutations may provide diagnostic clarity, though negative genetic testing does not definitively exclude HCM given incomplete understanding of disease genetics (Elliott et al., 2014).

CMR has emerged as the gold standard imaging modality for comprehensive cardiac evaluation in athletes with suspected pathological hypertrophy. CMR provides superior tissue characterization, precise volumetric measurements, and assessment of myocardial fibrosis through late gadolinium enhancement imaging (Kramer et al., 2008).

Standardized CMR protocols for athlete evaluation include assessment of biventricular volumes, function, and mass; evaluation of wall thickness and patterns of hypertrophy; tissue characterization to detect edema or fibrosis; and assessment of valvular function and flow dynamics (Kramer et al., 2008).

Reference ranges established from large population studies, such as the UK Biobank, provide context for interpreting findings in athletes, though sport-specific and ethnicity-specific variations must be considered (Petersen et al., 2017).

The presence of late gadolinium enhancement, particularly in non-ischemic patterns, raises significant concern for underlying cardiomyopathy and warrants comprehensive evaluation including consideration of myocardial biopsy in selected cases.

Comprehensive pre-participation cardiovascular screening of boxers should include detailed personal and family history, physical examination, and 12-lead ECG interpreted using athlete-specific criteria (Drezner et al., 2017). Athletes with abnormal findings require further evaluation with echocardiography and potentially CMR before medical clearance for competition.

Given the high cardiovascular demands of boxing and the potential for traumatic cardiac injury from chest blows, thorough screening is particularly important in this population.

Elite boxers benefit from periodic cardiovascular reassessment throughout their careers. Annual ECG screening and periodic echocardiographic evaluation enable tracking of cardiac adaptations and early detection of emerging pathology. Athletes demonstrating progressive wall thickness, chamber enlargement beyond expected physiological limits, or declining function warrant comprehensive investigation.

HRV monitoring represents a practical tool for continuous assessment of autonomic function and training adaptation that can be implemented at minimal cost using commercially available devices (Kiss et al., 2016; Morgan & Mora, 2017).

Athletes presenting with borderline findings that do not clearly distinguish physiological from pathological remodeling require individualized management. Multi-modality imaging including CMR, exercise stress testing, ambulatory ECG monitoring, and potentially cardiopulmonary exercise testing provide additional diagnostic information. Involvement of specialists in sports cardiology ensures optimal evaluation and management decisions.

In cases of persistent diagnostic uncertainty after comprehensive evaluation, shared decision-making with the athlete regarding continued competitive participation is essential, with full disclosure of known risks and regular reassessment.

For athletes who undergo detraining for diagnostic purposes, return to competition requires demonstration of regression of cardiac dimensions toward normal values, confirming the physiological nature of the original findings. Gradual return to training with periodic reassessment ensures safe resumption of competitive activity.

Despite significant advances in understanding cardiac adaptations in athletes, several areas warrant further investigation. Sport-specific reference ranges for cardiac parameters in boxers across different competitive levels, weight classes, and ethnic backgrounds would enhance diagnostic accuracy. Longitudinal studies tracking cardiac adaptations throughout boxing careers, including post-retirement follow-up, would provide insights into long-term consequences of intensive training.

Advanced imaging techniques including CMR feature tracking for strain analysis, perfusion imaging, and novel tissue characterization methods may further improve differentiation of physiological from pathological remodeling. Integration of HRV monitoring with other training load and recovery metrics could optimize individualized training prescription and enhance performance while minimizing health risks.

Finally, investigation of genetic and molecular markers of adaptive versus maladaptive cardiac remodeling may enable early identification of athletes at risk for adverse outcomes, allowing personalized preventive interventions.

Conclusion

Elite boxers develop characteristic cardiac structural adaptations in response to the unique physiological demands of their sport. These adaptations, including left ventricular hypertrophy, chamber enlargement, and altered autonomic regulation, represent normal physiological responses to training and contribute to enhanced athletic performance.

Distinguishing these physiological changes from pathological conditions such as hypertrophic cardiomyopathy remains a critical clinical challenge requiring comprehensive evaluation using established diagnostic criteria, advanced imaging modalities including CMR, and consideration of functional parameters including HRV. Gender-specific differences in adaptation patterns must be recognized when interpreting findings.

The application of athlete-specific ECG interpretation criteria, evidence-based imaging protocols, and novel assessment tools including HRV monitoring enables accurate diagnosis and appropriate management. Regular screening and longitudinal monitoring throughout athletic careers ensures early detection of potential pathology while allowing safe competitive participation for the vast majority of athletes with physiological adaptations.

Continued research into sport-specific cardiac adaptations, refinement of diagnostic criteria, and development of novel assessment techniques will further improve our ability to optimize both athlete safety and performance in boxing and other combat sports.

References

1. Bachynska, N.V., Doroshenko, E., Svatiev, A., & Bohuslavskyi, V. (2025). Assessing Gender Dimorphism Indicators in Anthropometric Characteristics and Features of the Bony Pelvis of Female Representatives of Pair and Group Types in Sports Acrobatics. *Physical Education Theory and Methodology*, *25*(1), 101-108. https://doi.org/10.17309/tmfv.2025.1.13

2. Bachynska N.V., Ivchenko O., Boguslavsky V., et al. (2023). Study of indicators of heart rate variability of acrobats at the stages of long-term training in the aspect of sexual dimorphism. *Acta Balneologica*, *65*, 3(175), 142–147.

3. Bianco A., Fischetti F., Cataldi S., Messina G., Thomas E. (2023). Physical Capacities and Combat Performance Characteristics of Male and Female

Olympic Boxers. *International Journal of Environmental Research and Public Health*, *20*(9), 5665. https://doi.org/10.3390/ijerph20095665

4. Chaabene H., Tabben M., Mkaouer B., et al. (2015). Amateur boxing: physical and physiological attributes. *Sports Medicine*, *45*(3), 337–352.

5. Davis P., Benson P.R., Pitty J.D., Connorton A.J., Waldock R. (2015). The activity profile of elite male amateur boxing. *International Journal of Sports Physiology and Performance*, *10*(1), 53–57.

6. Drezner J.A., Sharma S., Baggish A., et al. (2017). International criteria for electrocardiographic interpretation in athletes: Consensus statement. *British Journal of Sports Medicine*, *51*(9), 704–731.

7. Elliott P.M., Anastasakis A., Borger M.A., et al. (2014). 2014 ESC Guidelines on diagnosis and management of hypertrophic cardiomyopathy: the Task Force for the Diagnosis and Management of Hypertrophic Cardiomyopathy of the European Society of Cardiology (ESC). *European Heart Journal*, *35*(39), 2733–2779.

8. Ghani S., Sharma S. (2020). The athlete's heart. *British Journal of Hospital Medicine*, *81*(12), 1–9.

9. Harmon K.G., Asif I.M., Maleszewski J.J., et al. (2015). Incidence, cause, and comparative frequency of sudden cardiac death in National Collegiate Athletic Association athletes: a decade in review. *Circulation*, *132*(1), 10–19.

10. Kamandulis S., Venckunas T., Masiulis N., Snieckus A., Simonavicius J., Skurvydas A. (2018). Cardio-Respiratory Endurance Responses Following a Simulated 3 × 3 Minutes Amateur Boxing Contest in Elite Level Boxers. *Sports*, *6*(4), 119. https://doi.org/10.3390/sports6040119

11. Kiss O., Sydó N., Vargha P., Vágó H., Czimbalmos C., Édes E., et al. (2016). Detailed heart rate variability analysis in athletes. *Clinical Autonomic Research*, *26*(4), 245–252. https://doi.org/10.1007/s10286-016-0360-z

12. Kramer C.M., Barkhausen J., Flamm S.D., Kim R.J., Nagel E. (2008). Standardized cardiovascular magnetic resonance imaging (CMR) protocols, society for cardiovascular magnetic resonance: board of trustees task force on standardized protocols. *Journal of Cardiovascular Magnetic Resonance*, *10*, 35.

13. Leicht A.S., Halliday A., Sinclair W.H., D'Auria S., Buchheit M., Kenny G.P., Stanley J. (2018). Heart rate variability responses to acute and repeated postexercise sauna in trained cyclists. *Applied Physiology, Nutrition, and Metabolism*, *43*(7), 704–710. https://doi.org/10.1139/apnm-2017-0581

14. Lima-Borges D.S., Martinez P.F., Vanderlei L.C.M., Barbosa F.S.S., Oliveira-Junior S.A. (2018). Autonomic modulations of heart rate variability are associated with sports injury incidence in sprint swimmers. *The Physician and Sportsmedicine*, *46*(3), 374–384. https://doi.org/10.1080/00913847.2018.1450606

15. Morgan S.J., Mora J.A.M. (2017). Effect of Heart Rate Variability Biofeedback on Sport Performance, a Systematic Review. *Applied Psychophysiology and Biofeedback*, *42*(3), 235–245. https://doi.org/10.1007/s10484-017-9364-2

16. Petersen S.E., Aung N., Sanghvi M.M., et al. (2017). Reference ranges for cardiac structure and function using cardiovascular magnetic resonance (CMR) in Caucasians from the UK Biobank population cohort. *Journal of Cardiovascular Magnetic Resonance*, *19*(1), 18.

17. Plews D.J., Laursen P.B., Buchheit M. (2016). Day-to-day Heart Rate Variability (HRV) Recordings in World Champion Rowers: Appreciating Unique Athlete Characteristics. *International Journal of Sports Physiology and Performance*, *12*(5), 1–19. https://doi.org/10.1123/ijspp.2016-0343

18. Rabbani A., Clemente F.M., Kargarfard M., Chamari K. (2019). Match fatigue time-course assessment over four days: usefulness of the Hooper index and heart rate variability in professional soccer players. *Frontiers in Physiology*, *10*, 109. https://doi.org/10.3389/fphys.2019.00109

19. Riding N.R., Sheikh N., Adamuz C., et al. (2015). Comparison of three current sets of electrocardiographic interpretation criteria for use in screening athletes. *Heart*, *101*(5), 384–390.

20. Schneider C., Wiewelhove T., Raeder C., Flatt A.A., Hoos O., Hottenrott L., et al. (2019). Heart Rate Variability Monitoring During Strength and High-Intensity Interval Training Overload Microcycles. *Frontiers in Physiology*, *10*, 582. https://doi.org/10.3389/fphys.2019.00582

21. Utomi V., Oxborough D., Whyte G.P., et al. (2013). Systematic review and meta-analysis of training mode, imaging modality and body size influences on the morphology and function of the male athlete's heart. *Heart*, *99*(23), 1727–1733.