Petrushyn Dmytro, Cand. of Sciences Physical Education and Sports, Dnipro State University of Internal Affairs, Dnipro, Ukraine, https://orcid.org/0000-0001-9207-0804

CARDIOVASCULAR ADAPTATIONS IN ELITE BOXERS: A COMPARATIVE ANALYSIS WITH ENDURANCE ATHLETES

Abstract

Background: Different sports induce distinct patterns of cardiovascular adaptation based on their specific physiological demands. Boxing combines elements of both endurance and power training, creating a unique adaptive stimulus that differs from pure endurance sports. **Objective:** To compare and contrast cardiovascular adaptations in elite boxers with those observed in endurance athletes, examining structural remodeling, functional capacity, autonomic regulation, and metabolic characteristics. **Methods:** A comprehensive review of literature examining cardiac morphology, electrocardiographic patterns, heart rate variability, and physiological profiles in boxers, swimmers, rowers, and cyclists was conducted. Studies utilizing advanced imaging techniques and physiological assessments were analyzed to identify sport-specific adaptation patterns. **Results:** Elite boxers demonstrate a mixed pattern of cardiovascular adaptation, combining features of both endurance and resistance training adaptations. Compared to pure endurance athletes, boxers show more balanced ventricular hypertrophy with relatively less chamber dilation, different autonomic modulation patterns, and distinct metabolic profiles. Both groups demonstrate elevated heart rate variability compared to untrained individuals, though the magnitude and specific HRV parameters differ between sports. Gender-specific differences in adaptation patterns are evident across both boxing and endurance sports. **Conclusions:** Boxing induces unique cardiovascular adaptations that

reflect its combined endurance and power demands. Understanding these sport-specific differences is essential for appropriate clinical interpretation, training optimization, and development of tailored screening protocols. The comparative analysis provides insights into the plasticity of cardiovascular adaptation and the specificity of training responses.

Keywords: boxing, endurance athletes, cardiac remodeling, heart rate variability, athlete's heart, sports cardiology, comparative physiology, cardiovascular adaptation

Introduction

The human cardiovascular system demonstrates remarkable plasticity in response to chronic exercise training, with adaptations closely reflecting the specific physiological demands imposed by different sports (Utomi et al., 2013). While the concept of «athlete's heart» has been recognized for over a century, contemporary research reveals substantial heterogeneity in cardiovascular remodeling patterns across athletic disciplines.

Endurance sports such as swimming, cycling, and rowing have been extensively studied and serve as prototypical examples of volume-overload training that induces eccentric cardiac hypertrophy characterized by proportional increases in chamber dimensions and wall thickness (Utomi et al., 2013). In contrast, resistance training typically produces pressure-overload stimuli leading to more concentric hypertrophy patterns with greater wall thickening relative to chamber enlargement.

Boxing occupies a unique position in the spectrum of athletic disciplines, combining sustained aerobic efforts during training and competition with repeated explosive, high-intensity anaerobic bursts (Chaabene et al., 2015). The sport demands exceptional cardiovascular endurance to maintain high work rates

throughout multiple rounds, while simultaneously requiring explosive power for rapid punch combinations and defensive movements (Davis et al., 2015).

This mixed training stimulus creates cardiovascular adaptations that differ from pure endurance or resistance training, yet these sport-specific differences remain incompletely characterized. Understanding these distinctions has important implications for clinical evaluation of athletes, development of sport-specific screening protocols, training optimization, and assessment of cardiovascular health across athletic populations.

Recent advances in cardiac imaging, particularly cardiovascular magnetic resonance (CMR), refined electrocardiographic interpretation criteria for athletes, and widespread adoption of heart rate variability (HRV) monitoring have enabled more detailed characterization of sport-specific cardiovascular adaptations (Drezner et al., 2017; Kramer et al., 2008). Additionally, recognition of gender-specific differences in cardiovascular adaptation patterns has highlighted the need for sex-stratified analyses across different sports (Bianco et al., 2023).

This article provides a comprehensive comparative analysis of cardiovascular adaptations in elite boxers versus endurance athletes, examining structural remodeling, functional parameters, autonomic regulation, metabolic characteristics, and clinical implications of these sport-specific differences.

Elite boxers engage in multifaceted training regimens that typically include 6-12 sessions per week encompassing various modalities. Aerobic conditioning through road running or cycling forms the foundation, with sessions often lasting 30-60 minutes at moderate to high intensities. Technical training includes pad work, bag work, and sparring sessions that involve repeated high-intensity efforts lasting 2-3 minutes followed by brief rest periods (Chaabene et al., 2015).

During competitive bouts, boxers maintain heart rates between 85-95% of maximum throughout three-minute rounds, with activity profiles showing approximately 33% of time spent in high-intensity activity (Davis et al., 2015).

This creates a combined cardiovascular stimulus involving sustained elevated cardiac output with superimposed bursts of maximal effort.

Studies examining cardio-respiratory responses during simulated boxing contests demonstrate significant metabolic demands, with oxygen consumption reaching 70-85% of maximum and substantial anaerobic contributions during high-intensity exchanges (Kamandulis et al., 2018). The intermittent nature of boxing creates unique cardiovascular stress patterns distinct from the more steady-state demands of endurance sports.

Pure endurance athletes such as distance runners, cyclists, swimmers, and rowers typically train 10-20 hours per week primarily at moderate intensities with periodic high-intensity interval sessions. Training emphasizes sustained cardiovascular output over extended periods, with competitive events lasting from several minutes to several hours depending on the discipline.

For example, elite swimmers train 8-12 sessions weekly with daily distances often exceeding 10 kilometers, creating substantial volume-overload cardiovascular stimuli (Kamandulis et al., 2020). Similarly, world-class rowers engage in extensive training volumes combining on-water rowing with ergometer sessions and supplementary conditioning (Plews et al., 2016).

The cardiovascular workload in endurance sports predominantly involves volume overload with relatively modest pressure-overload components, contrasting with the mixed stimulus in boxing that includes both sustained cardiac output demands and explosive power requirements.

The metabolic profile of boxing differs substantially from pure endurance sports. While both require significant aerobic capacity, boxing involves greater reliance on anaerobic energy systems during high-intensity exchanges and punch combinations. Studies examining physiological profiles of Indian boxers reveal well-developed aerobic capacity combined with superior anaerobic power compared to general athletic populations (Singh et al., 2013).

Endurance athletes typically demonstrate higher maximal oxygen consumption (VO_2max) values than boxers, reflecting greater oxidative capacity developed through high training volumes. However, boxers show superior performance in tests of anaerobic power and repeated high-intensity efforts, adaptations specific to their sport's demands (Chaabene et al., 2015).

The pattern of left ventricular (LV) remodeling represents the most distinctive difference between boxers and endurance athletes. Endurance-trained athletes typically develop eccentric hypertrophy characterized by proportional increases in chamber dimensions and wall thickness, with LV end-diastolic volumes often exceeding 60-70 mL/m² and wall thickness generally remaining below 12 mm (Utomi et al., 2013).

In contrast, boxers demonstrate a more balanced remodeling pattern with moderate increases in both chamber dimensions and wall thickness, reflecting the combined volume and pressure overload of their training. Meta-analyses examining cardiac morphology across training modalities confirm that mixed-sport athletes, including boxers, show intermediate patterns between pure endurance and pure resistance training adaptations (Utomi et al., 2013).

The relative wall thickness (RWT), calculated as twice the posterior wall thickness divided by LV end-diastolic dimension, provides useful discrimination between remodeling patterns. Endurance athletes typically maintain normal or slightly reduced RWT due to proportional chamber and wall growth, while resistance-trained athletes may show elevated RWT. Boxers generally demonstrate RWT values between these extremes, reflecting their mixed training stimulus.

Reference ranges established from large population studies using standardized CMR protocols demonstrate that endurance athletes develop the most pronounced chamber enlargement among athletic populations (Petersen et al., 2017). LV end-diastolic volumes in elite endurance athletes frequently exceed

100-110 mL/m² in males and 80-90 mL/m² in females, representing substantial adaptations beyond normal population ranges.

Boxers show moderate chamber enlargement, typically with LV end-diastolic volumes of 90-100 mL/m² in males and 70-80 mL/m² in females, values intermediate between sedentary individuals and pure endurance athletes. This reflects adequate stroke volume enhancement for their sport's demands without the extreme chamber dilation seen in athletes training 15-25 hours weekly in endurance disciplines.

Right ventricular adaptation follows similar patterns, with endurance athletes demonstrating more pronounced RV enlargement compared to boxers. Standardized CMR protocols enable comprehensive assessment of biventricular remodeling across athletic populations (Kramer et al., 2008).

LV wall thickness measurements provide another point of differentiation between sports. Pure endurance athletes rarely develop wall thickness exceeding 12 mm, and values above 13 mm should prompt careful evaluation for pathology even in elite performers. The increased LV mass in endurance athletes results primarily from chamber dilation rather than wall thickening (Utomi et al., 2013).

Boxers may develop slightly greater wall thickness, with values of 12-13 mm being common in elite male competitors and occasional exceptional athletes approaching 14 mm while remaining within physiological limits (Chaabene et al., 2015). This greater relative wall thickness reflects the power and explosive strength components of boxing training that create pressure-overload stimuli.

Resistance-trained athletes, particularly those in strength sports, may develop even greater wall thickness (13-15 mm) with less chamber dilation, representing the most concentric pattern of athletic cardiac adaptation. Boxing thus occupies an intermediate position between endurance and strength sports in terms of hypertrophy patterns.

Left atrial enlargement occurs in both boxers and endurance athletes as an adaptation to enhance diastolic filling and stroke volume. Studies suggest that endurance athletes may develop more pronounced left atrial enlargement given the greater chronic volume overload their hearts experience during prolonged training sessions (Petersen et al., 2017).

However, excessive atrial enlargement in any athletic population warrants careful evaluation to exclude underlying pathology or assess potential arrhythmia risk, as marked atrial dilation may predispose to atrial fibrillation even in the absence of structural heart disease.

Recent research has increasingly recognized important gender differences in cardiovascular adaptation to athletic training across all sports disciplines. These differences have implications for establishing appropriate reference ranges and interpretation criteria.

Studies comparing male and female Olympic boxers reveal that while both sexes develop cardiac adaptations, males demonstrate more pronounced structural changes including greater absolute increases in LV mass, wall thickness, and chamber dimensions (Bianco et al., 2023). Female boxers show relatively greater improvements in diastolic function and contractile efficiency compared to structural remodeling magnitude.

Research examining gender dimorphism indicators in combat sports and acrobatic athletes demonstrates that physiological adaptations, including cardiovascular remodeling, follow sex-specific patterns related to hormonal influences, body composition differences, and potentially differential training responses (Bachynska et al., 2025). Heart rate variability patterns also differ between male and female combat athletes across training periods (Bachynska et al., 2023).

Similar gender patterns exist in endurance sports, with male athletes developing greater absolute cardiac dimensions while female athletes show larger

relative increases compared to sex-specific population norms. Studies in swimmers, for instance, demonstrate that while male swimmers develop greater absolute LV mass and chamber volumes, female swimmers show proportionally larger adaptations relative to baseline values (Kamandulis et al., 2020).

The clinical implication is that diagnostic criteria and reference ranges must account for sex-specific differences to avoid misclassification of physiological adaptations as pathological in male athletes or failure to detect pathology in female athletes.

Boxers commonly demonstrate training-related ECG changes including sinus bradycardia (often 40-50 bpm at rest), first-degree atrioventricular block, incomplete right bundle branch block, early repolarization patterns, and increased QRS voltage consistent with LV hypertrophy (Drezner et al., 2017). The prevalence and magnitude of these findings correlate with training volume and competitive level.

Application of international consensus criteria for ECG interpretation in athletes enables standardization of screening and reduces false-positive rates while maintaining sensitivity for detecting pathological conditions (Drezner et al., 2017). Comparative studies of different interpretation criteria demonstrate that athlete-specific approaches significantly improve specificity compared to standard clinical ECG criteria (Riding et al., 2015).

Endurance athletes typically demonstrate even more pronounced training-related ECG changes compared to boxers, reflecting their greater vagal tone and more pronounced cardiac remodeling. Sinus bradycardia may be more severe (30-45 bpm), and voltage criteria for LV hypertrophy are frequently met despite generally modest wall thickness increases, reflecting the chamber enlargement and enhanced stroke volume.

Early repolarization patterns, including J-point elevation and tall T waves, occur very commonly in endurance athletes and represent benign training

adaptations. However, certain patterns such as anterior T-wave inversions beyond V1, pathological Q waves, or QTc prolongation require investigation regardless of sport (Drezner et al., 2017).

Understanding sport-specific ECG patterns enables more accurate interpretation during pre-participation screening. While the basic principles of ECG evaluation remain consistent across sports, awareness that endurance athletes may demonstrate more pronounced changes helps avoid unnecessary investigations while maintaining appropriate vigilance for pathological findings.

The standardized approach outlined in international consensus statements provides a framework applicable across athletic populations while recognizing sport-specific variations (Drezner et al., 2017).

Heart rate variability analysis has emerged as a valuable tool for assessing autonomic function, training adaptation, and recovery status across athletic populations. Comparative analysis of HRV patterns between boxers and endurance athletes reveals both similarities and important differences.

Both boxers and endurance athletes demonstrate elevated HRV compared to sedentary populations, reflecting enhanced parasympathetic tone and cardiovascular autonomic regulation (Kiss et al., 2016). This adaptation contributes to resting bradycardia and represents a favorable training response associated with improved cardiac efficiency.

Detailed HRV analysis in athletes reveals complex patterns across multiple time-domain and frequency-domain parameters. Elite athletes across disciplines typically show increased RMSSD (root mean square of successive differences), SDNN (standard deviation of normal-to-normal intervals), and enhanced high-frequency power, all indicating robust parasympathetic activity (Kiss et al., 2016).

Studies comparing HRV across different sports suggest that endurance athletes may develop slightly greater parasympathetic dominance compared to mixed-sport athletes like boxers, reflecting their higher training volumes and

predominantly aerobic training stimulus. However, substantial individual variation exists, and sport-specific differences are generally smaller than the overall difference between trained and untrained individuals.

Research examining day-to-day HRV recordings in world champion rowers emphasizes the importance of appreciating individual athlete characteristics and establishing personalized baselines rather than relying solely on group comparisons (Plews et al., 2016). This principle applies across all sports, including boxing.

Studies in combat athletes including acrobats demonstrate that HRV adaptations parallel training progression and show gender-specific patterns, with male athletes typically demonstrating slightly greater absolute HRV values though both sexes show similar relative adaptations (Bachynska et al., 2023).

HRV monitoring provides practical applications for training load management across athletic populations. In endurance sports, HRV has been used to guide training intensity, with studies in swimmers and cyclists demonstrating associations between HRV patterns and injury risk, performance capacity, and optimal training prescription (Lima-Borges et al., 2018; Leicht et al., 2018).

For boxers and other combat athletes, HRV monitoring enables assessment of recovery from intensive training periods, detection of overreaching or overtraining, and optimization of tapering strategies before competition. Studies examining HRV during strength and high-intensity interval training overload microcycles demonstrate the utility of HRV for detecting excessive training stress (Schneider et al., 2019).

Professional team sport athletes, such as soccer players, benefit from HRV monitoring for assessing match fatigue and recovery time-course, principles applicable to boxers managing training loads and competition schedules (Rabbani et al., 2019).

Systematic reviews examining HRV biofeedback effects on sport performance suggest potential benefits across disciplines, though effects vary by sport type, athlete level, and intervention protocol (Morgan & Mora, 2017). HRV-guided training approaches show promise for optimizing performance while minimizing injury and illness risk.

For both boxers and endurance athletes, HRV assessment provides valuable insights into autonomic balance, training adaptation, and readiness to perform. Integration of HRV monitoring with other load and recovery metrics creates comprehensive athlete management systems.

Maximal oxygen consumption (VO_2max) represents the gold standard measure of aerobic capacity and differs substantially between boxers and endurance athletes. Elite endurance athletes in disciplines such as cross-country skiing, distance running, and cycling may achieve VO_2max values exceeding 70-80 mL/kg/min in males and 60-70 mL/kg/min in females, representing the highest values observed in athletic populations.

Elite boxers typically demonstrate VO_2max values of 55-65 mL/kg/min in males and 45-55 mL/kg/min in females, values substantially above general population norms but below pure endurance athletes (Chaabene et al., 2015; Singh et al., 2013). These values reflect adequate aerobic capacity for boxing's demands without the extreme oxidative adaptation of athletes training 15-25 hours weekly in endurance sports.

The lower relative VO_2max in boxers compared to endurance athletes does not indicate inadequate conditioning but rather sport-specific adaptation. Boxing success depends on multiple capacities beyond maximal aerobic power, including anaerobic capacity, power, speed, technical skill, and tactical acumen.

Boxers demonstrate superior anaerobic capacity and power compared to pure endurance athletes, reflecting their training emphasis on explosive movements and repeated high-intensity efforts. Studies examining physiological

profiles reveal that boxers excel in tests of anaerobic power, repeated sprint ability, and explosive strength compared to endurance-trained athletes (Chaabene et al., 2015).

This anaerobic emphasis creates different metabolic adaptations, with boxers showing greater buffering capacity, lactate tolerance, and phosphagen system development compared to endurance athletes who rely primarily on oxidative metabolism.

Both boxers and endurance athletes develop enhanced stroke volume compared to untrained individuals, though the magnitude differs. Endurance athletes with their greater chamber enlargement typically achieve larger absolute stroke volumes, often exceeding 100-120 mL at rest and 160-180 mL during maximal exercise in elite performers.

Boxers show moderate stroke volume enhancements adequate for their sport's demands, typically 80-100 mL at rest and 140-160 mL during maximal exercise. Combined with appropriate heart rate responses, this enables sufficient cardiac output for high-level boxing performance.

The maximal cardiac output during exercise may be greater in endurance athletes given both higher stroke volumes and ability to sustain elevated heart rates throughout prolonged efforts. However, boxers require adequate cardiac output during repeated high-intensity bursts rather than sustained maximal output, representing different cardiovascular demands.

Within combat sports, boxers share cardiovascular characteristics with other striking disciplines such as kickboxing and mixed martial arts, though differences exist based on specific sport demands. Studies examining judo athletes reveal similar mixed patterns of cardiac adaptation reflecting the combination of strength, power, and endurance requirements (Drid et al., 2015).

International versus national level comparisons within combat sports demonstrate that elite competitors develop more pronounced cardiovascular

adaptations, emphasizing the importance of competitive level when interpreting cardiac findings (Drid et al., 2015).

Team sport athletes such as soccer and basketball players demonstrate cardiovascular adaptations somewhat similar to boxers, reflecting the intermittent high-intensity nature of their sports. Studies examining heart rate variability in professional basketball players for assessing ventilatory thresholds reveal patterns intermediate between pure endurance and power athletes (Ramos-Campo et al., 2017).

However, the specific movement patterns, metabolic demands, and training structures differ substantially between team sports and boxing, creating nuanced differences in adaptation patterns.

The spectrum of cardiovascular adaptations across different sports demonstrates the remarkable plasticity and specificity of human physiological responses to training. Understanding this specificity enables:

1. More accurate clinical interpretation of cardiac findings in athletes from different disciplines
2. Recognition that reference ranges and diagnostic thresholds may require sport-specific adjustment
3. Appreciation that "athlete's heart" is not a single entity but rather a family of adaptations reflecting specific training stimuli
4. Development of sport-specific screening and monitoring protocols
5. Optimization of training approaches based on known adaptation patterns.

While comprehensive cardiovascular screening using standardized approaches applies across all sports (Drezner et al., 2017), awareness of sport-specific adaptation patterns improves interpretation accuracy. Clinicians evaluating boxers should recognize that modest wall thickness increases (12-13

mm) may be physiological even when slightly exceeding typical values for pure endurance athletes.

Conversely, the extreme chamber dilation sometimes observed in ultra-endurance athletes requires careful evaluation to ensure physiological adaptation rather than underlying cardiomyopathy, even when wall thickness remains normal.

Current diagnostic criteria for conditions such as hypertrophic cardiomyopathy utilize universal cutoff values (typically LV wall thickness ≥15 mm) regardless of athletic discipline (Elliott et al., 2014). While this approach maintains diagnostic sensitivity, awareness of sport-specific patterns helps clinicians assess borderline cases more accurately.

An endurance athlete with 14 mm wall thickness warrants more concern than a boxer with similar measurements, given that endurance training rarely produces such thickness. Conversely, marked chamber dilation may be more acceptable in endurance athletes than in boxers where such findings are less typical.

Understanding sport-specific cardiovascular adaptations enables more targeted training prescription. Boxers require sufficient endurance training to develop adequate aerobic capacity while maintaining focus on power, speed, and anaerobic capacity development. Excessive endurance training volume could potentially impair power and speed characteristics essential for boxing success.

Conversely, endurance athletes benefit from understanding that excessive concurrent strength training may interfere with optimal endurance adaptations, though some strength work enhances injury resistance and performance in most endurance disciplines.

Sport-specific demands influence return-to-sport decision-making after cardiac issues. The explosive, high-intensity nature of boxing creates substantial cardiovascular stress that may exceed safe limits for athletes with certain cardiac

conditions, even when those athletes might safely participate in moderate-intensity endurance activities.

Individualized assessment considering specific sport demands, cardiac condition severity, and athlete preferences enables optimal shared decision-making regarding competitive participation.

Comparing cardiovascular adaptations across sports presents methodological challenges including:

1. Selection of competitive level: Elite athletes in different sports may represent different percentiles of human performance capacity, complicating direct comparisons
2. Age and training history effects: Career length and age of training initiation vary across sports
3. Body composition differences: Sports select for different anthropometric characteristics that influence cardiac measurements
4. Imaging modality consistency: Standardized protocols are essential for valid comparisons
5. Gender representation: Many historical studies focused predominantly on male athletes, limiting understanding of female athletic adaptations.

The development and adoption of standardized cardiovascular assessment protocols across athletic populations enhances the validity of comparative research. Initiatives such as standardized CMR protocols and international consensus criteria for ECG interpretation represent important advances (Kramer et al., 2008; Drezner et al., 2017).

Future research should emphasize:

1. Longitudinal studies tracking athletes throughout their careers.
2. Inclusion of diverse athletic populations including underrepresented sports and demographic groups.

3. Multi-modality assessment combining imaging, electrocardiography, HRV analysis, and functional testing.

4. Genetic and molecular analyses to understand mechanisms underlying sport-specific adaptations.

5. Post-retirement follow-up to assess long-term cardiovascular health outcomes.

Practical Applications and Recommendations For Clinicians:

1. Recognize that «athlete's heart» encompasses diverse adaptation patterns reflecting sport-specific training stimuli.

2. Consider sport type when interpreting cardiac findings, using sport-specific expectations to guide clinical decision-making.

3. Apply standardized screening protocols while incorporating awareness of sport-specific variations.

4. Utilize multi-modality assessment including ECG, echocardiography, CMR, and HRV analysis for comprehensive evaluation.

5. Engage sports cardiology specialists for complex cases requiring differentiation of physiological from pathological findings.

For Coaches and Sports Scientists:

1. Understand cardiovascular demands of specific sports to optimize training prescription.

2. Implement HRV monitoring and other objective tools to assess training adaptation and recovery.

3. Balance training components appropriately for sport-specific demands (e.g., adequate endurance for boxing without excessive volume that impairs power development).

4. Recognize individual variation in adaptation rates and patterns, personalizing training approaches accordingly.

5. Collaborate with medical professionals to ensure athlete cardiovascular health while optimizing performance.

For Athletes:

1. Understand that cardiovascular adaptations reflect normal responses to training and contribute to performance.

2. Participate in recommended cardiovascular screening to ensure structural and electrical cardiac health.

3. Utilize available monitoring tools (HRV, subjective wellness measures) to optimize training and recovery.

4. Communicate with coaches and medical professionals about unusual symptoms or concerns.

5. Maintain perspective that while athletic training produces cardiac changes, these generally represent healthy adaptations.

Conclusion

Cardiovascular adaptations in elite boxers differ substantially from those observed in pure endurance athletes, reflecting the unique physiological demands of boxing that combine endurance, power, speed, and repeated high-intensity efforts. Boxers demonstrate a mixed pattern of cardiac remodeling intermediate between the eccentric hypertrophy of endurance athletes and the more concentric patterns of strength athletes, with moderate chamber enlargement and balanced wall thickness increases.

Both boxers and endurance athletes develop enhanced parasympathetic tone reflected in elevated heart rate variability, though endurance athletes may show slightly greater magnitude given their higher training volumes. Gender-specific differences in adaptation patterns exist across both sports, with males generally demonstrating more pronounced structural changes while females show relatively greater functional improvements.

Understanding these sport-specific differences has important clinical implications for cardiovascular screening, interpretation of cardiac findings, differentiation of physiological from pathological conditions, and optimization of training approaches. The application of standardized assessment protocols combined with awareness of sport-specific adaptation patterns enables accurate diagnosis and appropriate management.

Comparative analysis of cardiovascular adaptations across sports demonstrates the remarkable plasticity of the cardiovascular system and the specificity of training responses. This understanding enhances both clinical care of athletes and optimization of training methodologies. Continued research examining sport-specific adaptations, particularly longitudinal studies and investigations including diverse and underrepresented populations, will further refine our understanding and improve care for athletes across all disciplines.

The integration of advanced imaging techniques, HRV monitoring, genetic analyses, and functional assessments promises to provide increasingly sophisticated understanding of cardiovascular adaptation mechanisms and enable truly personalized approaches to athlete assessment, training, and medical care. As sports science and sports cardiology continue evolving, the goal remains optimizing both athlete performance and long-term cardiovascular health across the diverse spectrum of athletic disciplines.

References

1. Bachynska, N.V., Doroshenko, E., Svatiev, A., & Bohuslavskyi, V. (2025). Assessing Gender Dimorphism Indicators in Anthropometric Characteristics and Features of the Bony Pelvis of Female Representatives of Pair and Group Types in Sports Acrobatics. *Physical Education Theory and Methodology*, *25*(1), 101-108. https://doi.org/10.17309/tmfv.2025.1.13

2. Bachynska N.V., Ivchenko O., Boguslavsky V., et al. (2023). Study of indicators of heart rate variability of acrobats at the stages of long-term training in the aspect of sexual dimorphism. *Acta Balneologica*, *65*, 3(175), 142–147.

3. Bianco A., Fischetti F., Cataldi S., Messina G., Thomas E. (2023). Physical Capacities and Combat Performance Characteristics of Male and Female Olympic Boxers. *International Journal of Environmental Research and Public Health*, *20*(9), 5665. https://doi.org/10.3390/ijerph20095665

4. Chaabene H., Tabben M., Mkaouer B., et al. (2015). Amateur boxing: physical and physiological attributes. *Sports Medicine*, *45*(3), 337–352.

5. Davis P., Benson P.R., Pitty J.D., Connorton A.J., Waldock R. (2015). The activity profile of elite male amateur boxing. *International Journal of Sports Physiology and Performance*, *10*(1), 53–57.

6. Drezner J.A., Sharma S., Baggish A., et al. (2017). International criteria for electrocardiographic interpretation in athletes: Consensus statement. *British Journal of Sports Medicine*, *51*(9), 704–731.

7. Drid P., Casals C., Mekić A., Radjo I., Stojanović M., Ostojić S.M. (2015). Fitness and anthropometric profiles of international vs. national judo medalists in half-heavyweight category. *Journal of Strength and Conditioning Research*, *29*(8), 2115–2121.

8. Elliott P.M., Anastasakis A., Borger M.A., et al. (2014). 2014 ESC Guidelines on diagnosis and management of hypertrophic cardiomyopathy: the Task Force for the Diagnosis and Management of Hypertrophic Cardiomyopathy of the European Society of Cardiology (ESC). *European Heart Journal*, *35*(39), 2733–2779.

9. Ghani S., Sharma S. (2020). The athlete's heart. *British Journal of Hospital Medicine*, *81*(12), 1–9.

10. Kamandulis S., Juodsnukis A., Stanislovaitiene J., Zuoziene I.J., Bogdelis A., Mickevicius M., et al. (2020). Daily Resting Heart Rate Variability in

Adolescent Swimmers during 11 Weeks of Training. *International Journal of Environmental Research and Public Health*, *17*(6), E2097. https://doi.org/10.3390/ijerph17062097

11. Kamandulis S., Venckunas T., Masiulis N., Snieckus A., Simonavicius J., Skurvydas A. (2018). Cardio-Respiratory Endurance Responses Following a Simulated 3 × 3 Minutes Amateur Boxing Contest in Elite Level Boxers. *Sports*, *6*(4), 119. https://doi.org/10.3390/sports6040119

12. Kiss O., Sydó N., Vargha P., Vágó H., Czimbalmos C., Édes E., et al. (2016). Detailed heart rate variability analysis in athletes. *Clinical Autonomic Research*, *26*(4), 245–252. https://doi.org/10.1007/s10286-016-0360-z

13. Kramer C.M., Barkhausen J., Flamm S.D., Kim R.J., Nagel E. (2008). Standardized cardiovascular magnetic resonance imaging (CMR) protocols, society for cardiovascular magnetic resonance: board of trustees task force on standardized protocols. *Journal of Cardiovascular Magnetic Resonance*, *10*, 35.